arXiv:0905.3029v1 [math.GN] 19 May 2009

A NOTE ON THE COMMUTATIVITY OF INVERSE LIMIT AND ORBIT MAP

MAHENDER SINGH

Abstract. We show that the inverse limit and the orbit map commute for actions of compact groups on compact Hausdorff spaces.

1. Introduction

This note is motivated by the following example of Bredon [1, p.145]. Let S^2 be the 2-sphere identified with the unreduced suspension of the circle $S^1 = \{z \in \mathbb{C}\ ;\ |z| = 1\}$, and $f : S^2 \to S^2$ be the suspension of the map $S^1 \to S^1$, $z \mapsto z^3$. Then f commutes with the antipodal involution on S^2. If Σ is the inverse limit of the inverse system

$$\cdots \xrightarrow{f} S^2 \xrightarrow{f} S^2 \xrightarrow{f} S^2$$

then $\Sigma/\mathbb{Z}_2$ is homeomorphic to $\varprojlim \mathbb{R}P^2$.

We show that this can be generalized, that is, the inverse limit and the orbit map commute for actions of compact groups on compact Hausdorff spaces. The proof of the result is simple, but does not seem to be available in the literature. Before we prove the result, we recall basic notions that will be used throughout the note. By a map we mean a continuous function of topological spaces.

An inverse system of topological spaces, denoted by $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$, consists of a directed set Λ, a family of topological spaces $\{X_\alpha\}_{\alpha\in\Lambda}$ and a collection of maps $\pi_\alpha^\beta : X_\beta \to X_\alpha$ for $\alpha < \beta$ in Λ satisfying:

(1) $\pi_\alpha^\alpha : X_\alpha \to X_\alpha$ is the identity map for all $\alpha \in \Lambda$;
(2) $\pi_\alpha^\beta \pi_\beta^\gamma = \pi_\alpha^\gamma$ for all $\alpha < \beta < \gamma$ in Λ.

The maps π_α^β are called bonding maps. Given an inverse system $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$ of topological spaces, let $\varprojlim X_\alpha$ (possibly empty) be the subset of $\Pi_{\alpha\in\Lambda} X_\alpha$ consisting of elements (x_α) such that $x_\alpha = \pi_\alpha^\beta(x_\beta)$ for $\alpha < \beta$. It is given the subspace topology from the product space $\Pi_{\alpha\in\Lambda} X_\alpha$ and is called the inverse limit of the inverse system $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$. We denote by $\pi_\beta : \varprojlim X_\alpha \to X_\beta$, the restriction of the canonical projection $\Pi_{\alpha\in\Lambda} X_\alpha \to X_\beta$. The following basic results are well known.

Date: April 2009.
2000 Mathematics Subject Classification. Primary 54B25; Secondary 57S10.
Key words and phrases. Equivariant map; group action; inverse system.

Theorem 1.1. [2, p.217] *The inverse limit of an inverse system of non-empty compact Hausdorff topological spaces is a non-empty compact Hausdorff topological space.*

Theorem 1.2. [2, p.219] *Let X be a topological space and $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$ be an inverse system of topological spaces. If for each $\alpha \in \Lambda$ there is a map $\psi_\alpha : X \to X_\alpha$ such that $\pi_\alpha^\beta \psi_\beta = \psi_\alpha$ for $\alpha < \beta$, then there is a unique map $\psi : X \to \varprojlim X_\alpha$.*

We refer the reader to [2] for more details on inverse systems.

If $\{G_\alpha, \nu_\alpha^\beta, \Lambda\}$ is an inverse system of topological groups (ν_α^β are topological group homomorphisms), then $\varprojlim G_\alpha$ is also a topological group with group operation $\big((g_\alpha), (h_\alpha)\big) \mapsto (g_\alpha h_\alpha)$.

An action of a topological group G on a topological space X is a map $G \times X \to X$ denoted by $(g, x) \mapsto g \cdot x$ satisfying the following:

(1) $e \cdot x = x$ for all $x \in X$, where $e \in G$ is the identity element;
(2) $g \cdot (h \cdot x) = gh \cdot x$ for all $g,\ h \in G$ and $x \in X$.

We say that X is a G-space if there is an action of G on X. For each $x \in X$, $\overline{x} = \{g \cdot x \mid g \in G\}$ is called the orbit of x. There is associated with a G-space X, the orbit space X/G of all orbits. The canonical map $X \to X/G$ given by $x \mapsto \overline{x}$ is called the orbit map. We say that G acts freely on X if for each $x \in X$, $g \cdot x = x$ implies that $g = e$. We will use the following basic result.

Theorem 1.3. [1, p.38] *If G is a compact topological group acting on a compact Hausdorff topological space X, then X/G is also a compact Hausdorff topological space.*

Let X be a G-space, Y be a H-space and $\nu : G \to H$ be a topological group homomorphism. Then a map $f : X \to Y$ is called ν-equivariant if

$$f(g \cdot x) = \nu(g) \cdot f(x) \text{ for all } g \in G \text{ and } x \in X.$$

If both X and Y are G-spaces, then f is called G-equivariant. Note that the ν-equivariant map f induces a map $\overline{f} : X/G \to Y/H$ given by $\overline{f}(\overline{x}) = \overline{f(x)}$. If $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$ is an inverse system of topological spaces and $\{G_\alpha, \nu_\alpha^\beta, \Lambda\}$ is an inverse system of topological groups where each X_α is a G_α-space and each bonding map π_α^β is ν_α^β-equivariant, then we get another inverse system $\{X_\alpha/G_\alpha, \overline{\pi_\alpha^\beta}, \Lambda\}$ by passing to orbit spaces. Also, under above conditions $\varprojlim X_\alpha$ is a $\varprojlim G_\alpha$-space with the action given by

$$(g_\alpha) \cdot (x_\alpha) = (g_\alpha \cdot x_\alpha) \text{ for } (g_\alpha) \in \varprojlim G_\alpha \text{ and } (x_\alpha) \in \varprojlim X_\alpha.$$

The reader can consult [1] for more details on group actions.

2. Commutativity of inverse limit and orbit map

In view of the above discussion it is natural to ask, when is $(\varprojlim X_\alpha)/(\varprojlim G_\alpha)$ homeomorphic to $\varprojlim(X_\alpha/G_\alpha)$. We present the following theorem in this direction.

Theorem 2.1. *Let $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$ be an inverse system of non-empty compact Hausdorff topological spaces and let $\{G_\alpha, \nu_\alpha^\beta, \Lambda\}$ be an inverse system of compact topological groups, where each X_α is a G_α-space and each bonding map π_α^β is ν_α^β-equivariant. Further, assume that Λ has the least element λ, G_λ action on X_λ is free and the bonding map ν_λ^α is injective for each $\alpha \in \Lambda$. Then there is a homeomorphism*

$$\psi : (\varprojlim X_\alpha)/(\varprojlim G_\alpha) \to \varprojlim(X_\alpha/G_\alpha).$$

We first prove the following simple lemma.

Lemma 2.2. *Let $\{X_\alpha, \pi_\alpha^\beta, \Lambda\}$ be an inverse system of non-empty compact Hausdorff topological spaces and let $\{G_\alpha, \nu_\alpha^\beta, \Lambda\}$ be an inverse system of compact topological groups, where each X_α is a G_α -space and each bonding map π_α^β is ν_α^β-equivariant. Then there is a closed continuous surjection*

$$\psi : (\varprojlim X_\alpha)/(\varprojlim G_\alpha) \to \varprojlim(X_\alpha/G_\alpha).$$

Proof. Let $X = \varprojlim X_\alpha$ and $G = \varprojlim G_\alpha$. Let $\pi_\beta : X \to X_\beta$ and $\nu_\beta : G \to G_\beta$ be the canonical projections for each $\beta \in \Lambda$. Clearly each π_β is ν_β-equivariant and therefore induces a map $\psi_\beta : X/G \to X_\beta/G_\beta$ given by $\overline{(x_\alpha)} \mapsto \overline{x_\beta}$ (note that $\psi_\beta = \overline{\pi_\beta}$). Also observe that for $\gamma < \beta$ the diagram

$$\begin{tikzcd} X/G \arrow[r, "\psi_\beta"] \arrow[d, "\psi_\gamma"] & X_\beta/G_\beta \arrow[dl, "\overline{\pi_\gamma^\beta}"] \\ X_\gamma/G_\gamma & \end{tikzcd}$$

commutes. Therefore by Theorem 1.2, we have the map

$$\psi : X/G \to \varprojlim(X_\alpha/G_\alpha)$$

given by $\overline{(x_\alpha)} \mapsto (\overline{x_\alpha})$. Clearly ψ is surjective. By Theorem 1.1 and Theorem 1.3, we see that X/G is compact and $\varprojlim(X_\alpha/G_\alpha)$ is Hausdorff. Therefore ψ is a closed map. This completes the proof of the lemma. □

We now complete the proof of Theorem 2.1.

Proof. It just remains to show that the map ψ is injective. Suppose $(\overline{x_\alpha}) = (\overline{y_\alpha})$. This means $\overline{x_\alpha} = \overline{y_\alpha}$ for each $\alpha \in \Lambda$. Thus for $\alpha \in \Lambda$, we have $x_\alpha = g_\alpha \cdot y_\alpha$ and $x_\lambda = g_\lambda \cdot y_\lambda$ for some $g_\alpha \in G_\alpha$ and $g_\lambda \in G_\lambda$. Since $\lambda \in \Lambda$ is the least element, we have $\lambda < \alpha$ and hence $\pi_\lambda^\alpha(x_\alpha) = x_\lambda$ and $\pi_\lambda^\alpha(y_\alpha) = y_\lambda$. This gives $x_\lambda = \pi_\lambda^\alpha(x_\alpha) = \pi_\lambda^\alpha(g_\alpha \cdot y_\alpha) = \nu_\lambda^\alpha(g_\alpha) \cdot \pi_\lambda^\alpha(y_\alpha) = \nu_\lambda^\alpha(g_\alpha) \cdot y_\lambda$ and hence $g_\lambda \cdot y_\lambda = \nu_\lambda^\alpha(g_\alpha) \cdot y_\lambda$. The freeness of G_λ action on X_λ implies $g_\lambda = \nu_\lambda^\alpha(g_\alpha)$.

Now, for $\lambda < \alpha < \beta$ we have $\nu_\lambda^\alpha \nu_\alpha^\beta(g_\beta) = \nu_\lambda^\beta(g_\beta) = g_\lambda$. By injectivity of ν_λ^α we get $\nu_\alpha^\beta(g_\beta) = g_\alpha$. Thus $(g_\alpha) \in G$ and $(x_\alpha) = (g_\alpha \cdot y_\alpha) = (g_\alpha) \cdot (y_\alpha)$. This shows that ψ is injective and hence a homeomorphism by Lemma 2.2. $\square$

As a consequence we have the following corollary.

Corollary 2.3. *Let G be a compact topological group acting freely on a compact Hausdorff topological space X and let $f : X \to X$ be a G-equivariant map. Then for the inverse system*

$$\cdots \xrightarrow{f} X \xrightarrow{f} X \xrightarrow{f} X$$

$(\varprojlim X)/G$ is homeomorphic to $\varprojlim(X/G)$.

References

1. G.E.Bredon, *Introduction to Compact Transformation Groups*, Academic Press, New York, 1972.
2. S.Eilenberg, N.E. Steenrod, *Foundations of Algebraic Topology*, Princeton University Press, 1952.

School of Mathematics, Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad-211019, INDIA.

E-mail address: `msingh@mri.ernet.in`